2.B: 2.N

TRANSFER REACTIONS IN THE INTERACTION OF ⁴⁰Ar WITH ²³²Th

A. G. ARTUKH, G. F. GRIDNEV, V. L. MIKHEEV, V. V. VOLKOV and J. WILCZYNSKI[†] Joint Institute for Nuclear Research, Dubna, USSR

Received 13 April 1973

Abstract: Transfer reactions in the bombardment of a ²³²Th target with 297 MeV and 388 MeV ⁴⁰Ar ions have been studied. The energy spectra, angular distributions and the cross sections for the formation of products having atomic numbers $5 \le Z \le 20$ have been measured. In the energy spectra of the reaction products broad peaks having energies in the vicinity of the output Coulomb barrier height have been observed along with the quasi-elastic ones. The dependence of the partial angular distribution of the light products on the reaction *Q*-value has been determined. The observed partial angular distributions have been analysed by applying the Strutinsky model. The total cross sections of light product formation have been found to be 1100 mb and 2400 mb for the ⁴⁰Ar energies E = 297 MeV and E = 388 MeV, which give the values of the critical angular momentum of 102 \hbar and 94 \hbar respectively for the two cases.

Е

NUCLEAR REACTIONS ²³²Th(⁴⁰Ar, X), E = 297, 388 MeV; measured $\sigma(E_X, \vartheta)$ for X = B, C, N, O, F, Ne, Na, Mg, Al, Si, P, S, Cl, Ar, K, Ca. Natural target.

1. Introduction

Nuclear reactions induced by the heavy ions such as Ar, Zn, Kr, Xe [refs. $^{1-4}$)], have become of great importance in the recent years. Reactions induced by such ions may result in the synthesis of superheavy elements in a new stability region. Transfer reactions in the bombardment of heavy nuclei by 40 Ar ions proved to be an effective means for obtaining new neutron-rich isotopes of light elements ⁵). Therefore, the study of the peculiarities of the mechanism of nuclear reactions with heavy ions and the yield of various products, both light and heavy, becomes still more significant.

The majority of investigations of direct heavy-ion reactions on heavy nuclei have been performed by radiochemical separation of reaction products ^{1, 6-8}). However, this method does not allow the detection of stable products as well as radioactive products having either very short or very long half-lives. It is even more difficult to measure their energy spectra. The combination of the magnetic analysis and the ΔE , E methods ⁹) is free from such disadvantages and makes it possible to identify unambiguously the light products of reactions over a wide range of A and Z. However, this technique used for measuring the angular and energy distributions of many reaction products requires much accelerator time. The ΔE , E method is more efficient, though it involves some difficulties in isotope separation, especially with Z > 8[refs. ^{10, 11}]].

[†] On leave from the Institute of Nuclear Physics, Crakow, Poland.

A. G. ARTUKH et al.

The study of some general features of the direct reactions such as the total cross sections and the cross sections for the production of some element does not require the knowledge of the mass number of products when reliable Z separation has been done. Some reaction properties such as the angular distributions of neighbouring isotopes differ slightly ^{6, 8, 12}). Earlier, in bombarding ²³²Th with ⁴⁰Ar ions we have found that the major contribution (of 75–85%) to the formation cross section of elements at an angle of 40° comes from 3–4 neighbouring isotopes ⁵). Angular and energy distributions for the given element may be expected to be close to those of separate isotopes which form the major part of the element produced.

The present investigation was aimed at measuring the cross sections, angular and energy distributions of the light reaction products formed in the bombardment of 232 Th by 297 MeV and 388 MeV 40 Ar ions using the "element" approach.

2. Experimental procedure

The ${}^{40}Ar(+7)$ and ${}^{40}Ar(+8)$ external beams of 2×10^{11} and 6×10^{10} particles/sec and energies 297 MeV and 388 MeV, respectively, from the 310 cm Dubna cyclotron were used for bombarding a metallic 232 Th target.

Reaction products were detected by a telescope consisting of two silicon surfacebarrier detectors: ΔE detector 27 μ m thick and the total absorption detector (E). The aperture was defined by a diaphragm of 5 mm in diameter placed in front of the ΔE detector at 250 mm from the target. The angular resolution of the telescope was found to be not worse than $\pm 1.5^{\circ}$, with the beam spot dimension of 6×6 mm² and the angular divergence of the beam less than $\pm 0.5^{\circ}$. The relative accuracy of the telescope location was $\pm 0.1^{\circ}$. A thin gold foil, 130 μ g/cm² thick, protected the detectors from the slow electron flux emitted by the bombarded target. Pulses from both the detectors were fed into a converter for a two-parameter pulse-height analysis. The twodimensional ΔE , $(E - \Delta E)$ spectrum was recorded in two 4096-channel analysers operating in the 256 $(\Delta E) \times 32(E - \Delta E)$ channel mode. Product yields at various angles were normalized with respect to the number of elastically scattered ⁴⁰Ar ions detected in a monitor solid state detector which was placed at an angle of 30° to the beam.

3. Experimental results

3.1. ENERGY SPECTRA

Since the energy losses of detected ions in a target were considerable, it was assumed that nuclear reactions occur in a middle plane of the target. The energy losses of reaction products in a target were determined by comparing the energy of 40 Ar ions elastically scattered from a 197 Au target of thickness 130 µg/cm² and the 232 Th target used in our experiment. By employing d*E*/dx tables for heavy ions (in Au and U) of Northcliffe and Shilling 13) it was found that our 232 Th target is equivalent to 3.9 mg/cm² of uranium. This value together with the data on dE/dx was later used to determine the energy losses of products in the target. Earlier, in bombarding ²³²Th by ⁴⁰Ar ions of energy 290 MeV it was established that at an angle of 40° the following isotopes: ¹⁹O, ²¹F, ²⁴Ne, ²⁶Na, ²⁸Mg, ³⁰Al, ³³Si, ³⁵P, ³⁶S, ³⁹Cl, ⁴⁰Ar,



ENERGY (MeV)

Fig. 1. The laboratory energy spectra of light products obtained in the bombardment of ²³²Th with 297 MeV ⁴⁰Ar ions. The product energies have been recalculated for energies in the middle plane of the target. The curves have been drawn through experimental points. All the curves and points for each detection angle have been multiplied by the coefficients in the top right-hand side of the figure.

⁴²K and ⁴⁴Ca [ref. ⁵)] have the maximum yield. All the calculations were made for the mass numbers of the above isotopes. It is worth noting that these isotopes are obtained with the total number of transferred nucleons close to twice the number of transferred protons. Such a ratio has been obtained by the authors for the 232 Th+ 22 Ne reaction 14).

The absolute values of product yields were calculated from the ratio of reaction product yields and elastically scattered ⁴⁰Ar at angles of 20–30° for 297 MeV and 20–25° for 388 MeV. It is possible to consider the $\sigma_{\rm el}/\sigma_{\rm Ruth}$ ratio to be unity for these angles ¹⁵).

A. G. ARTUKH et al.

Figs. 1 and 2 show the energy spectra of different products from Mg to Ca in the lab. system, the ⁴⁰Ar energy in the middle plane of the target being $E_0 = 288$ MeV. The energy spectra of lighter products and all the products at larger detection angles are not given since the low yield of these products does not make it possible to plot them with sufficient accuracy. However an evaluation of their formation cross section was made and this was used in plotting their angular distributions.



Fig. 2. See caption to fig. 1.

Figs. 3, 4 and 5 show the energy spectra of different products from O to Ca at the ⁴⁰Ar ion energy of $E_0 = 379$ MeV. Some properties characteristic of the energy spectra are worth noting.

The spectra cover rather a wide energy region: 80–120 MeV for $E_0 = 288$ MeV and 110–220 MeV for $E_0 = 379$ MeV. A comparison of the energy spectra of elements and their isotopes emitted at the angle of 40° in irradiation of a ²³²Th target with ²²Ne ions of 174 MeV and ⁴⁰Ar ions of 290 MeV show that the widths of the element and isotope spectra are close ^{5, 14}). Thus, one may consider that the energy spread introduced into the element energy spectrum by summing up the spectra of separate

isotopes is considerably smaller than the dynamic dispersion of the product energy due to the mechanism of interaction of two nuclei in direct reactions.

The energy spectra of the products in multi-nucleon transfer reactions usually have an approximately bell-shaped form. With the lowering of the number of transferred protons, the maxima in the energy spectra are shifted towards higher energies, while the spectra themselves become more asymmetric.



ENERGY (MeV)

Fig. 3. The laboratory energy spectra of light products obtained in the bombardment of ²³²Th with 388 MeV ⁴⁰Ar ions. The product energies have been recalculated for the energies in the middle plane of the target. The curves have been drawn through experimental points. All the curves and points for each detection angle have been multiplied by the coefficients shown in the top right-hand side of the figure.

The energy of incident particles greatly affects the widths and shapes of energy spectra. Thus, in increasing energy from 288 to 379 MeV the FWHM for Mg are changed from 40-50 MeV to 80-90 MeV. The effect of incident particle energy on the shape of the energy spectra is shown in fig. 6 where the P, Cl and K energy spectra are compared for various detection angles and the incident particle energies. The arrows in fig. 6 indicate the energy of the light products in the lab. system calculated under the assumption that the nuclear system decaying into two products does so when the kinetic energy of relative motion is zero (i.e. corresponding to the Coulomb barrier height for spherical nuclei). It should be noted that if we take into account







ENERGY (MeV) Fig. 5. See caption to fig. 3.

the velocity of the centre of mass the low-energy cut-offs of the spectra for the two incident energies are really equal.

At the ⁴⁰Ar energy of $E_0 = 379$ MeV the spectra of products of few-nucleon transfer reactions at the angles of 35° - 45° show two broad peaks one at a low-energy and the other at a high energy (fig. 5). For $E_0 = 288$ MeV such a spectra separation into two parts at large angles is observed for K and Ca only (fig. 2).

In general, for few-nucleon transfer reactions, the energy spectra of products tend to become softer with decreasing angle of emission.





Fig. 6. The dependence of the energy spectrum of the light product on the incident ion energy. The open circles are the 40 Ar energy E = 388 MeV, the filled ones are E = 297 MeV. All the curves and points for each detection angle are multiplied by the coefficients shown in the top right-hand side of the figure. The arrows show product energies for the system decaying into two spherical nuclei from the Coulomb barrier height. The dashed arrows are drawn for the 40 Ar energy E = 388 MeV, the solid arrows are drawn for E = 297 MeV.

3.2. ANGULAR DISTRIBUTIONS

The differential cross sections $d\sigma(\vartheta)/d\Omega$ of light product formation for two values of ⁴⁰Ar ion energy are shown in fig. 7. The cross sections have been obtained by integrating the energy spectra $d^2\sigma(\vartheta, E)/dE \cdot d\Omega$ over energy. Neutron transfer channels and ⁴⁰Ar inelastic scattering were separated by subtracting the normalized elastic scattering peak for small angles (10-15°) from the Ar energy spectrum at an angle ϑ . Only a part of the spectrum located 10 MeV below the calculated maximum of the elastic peak at $E_0 = 288$ MeV and 7 MeV below at $E_0 = 379$ MeV safely separated over the whole angular range. It is this part of the spectrum that was integrated when obtaining Ar angular distribution. Note that in such a treatment the contribution of the Coulomb excitation to the cross section is practically excluded ¹⁶).

As is seen from fig. 7 the shape of angular distributions varies in a regular fashion with the number of nucleons transferred. The peculiar peaks at somewhat smaller angles than that of the ⁴⁰Ar Rutherford scattering in the grazing collision is clearly seen only for few-nucleon transfer reactions. (The angles of the ⁴⁰Ar Rutherford



Fig. 7. The laboratory angular distributions of light products for 40 Ar ion energies in the middle plane of the target $E_0 = 288$ MeV and $E_0 = 379$ MeV. Solid curves are drawn throught the experimental points. All the points and curves for each element have been multiplied by the coefficients shown in the figure.

TABLE 1 Pross section of light product formation (in mb) in the ²³² Th+ ⁴⁰ Ar (297 MeV and 388 MeV) reactions
--

Energy ⁴⁰ Ar <i>E</i> (MeV)	Product	B	C	z	0	Ľ.	Ne	Na	Mg	A	Si	P 4	s	G	Ar ^a)	×	Ca	$\Sigma_i Z_i$
297	Jmcas						3.8	4.4	8.9	18	39	65	150	220	390	52	24	970
	£						8	6	16	25	50	78	170	240	410	65	38	1100
388	Gmeas	10	16	Π	13	Ξ	18	24	41	46	78	110	210	750	650	110	48	1700
	α	18	27	19	26	21	33	43	65	86	150	190	300	380	750	140	140	2400

^a) Ar yield has been determined by using a part of the energy spectrum, see subsect. 3.2 of the text.

scattering in the lab. coordinate system are 55° and 36°, for $E_0 = 288$ MeV and $E_0 = 379$ MeV respectively, and $r_0 = 1.46 \times 10^{-13}$ cm.)

With increasing number of transferred nucleons the width of these peaks of the angular distributions is increased, while its maximum is shifted towards smaller angles. In multinucleon transfer reactions the differential cross sections of products are increased monotonically with decreasing emission angle. For the reactions with the transfer of the maximum number of nucleons investigated here (C, N, O, for $E_0 = 379$ MeV) one observes some increase of product yields at larger detection angles. The increase of cross section of an angle of 10° for C, N, O, Ca requires a more detailed study since it can be connected with the background effects, e.g., of C and O admixtures in the target.

In earlier investigations $^{6, 8, 17}$) on the angular distributions of the products of transfer reactions on heavy nuclei both these features of maxima at the Rutherford scattering angles and of monotonically increasing yield towards 0° have been obtained. However, only a small number of reaction channels have been detected in the above studies. So it was difficult to determine a definite dependence of the angular distribution shape upon the number of transferred nucleons. The data shown in fig. 7, with the transfer up to about thirty nucleons, allow one to follow the evolution of the angular distribution shape with the increase in the number of transferred nucleons. A similar result for the angular distributions of direct reaction products (light elements from Li to Mg) was obtained by the authors 14) in the bombardment of 232 Th with 22 Ne ions.

3.3. LIGHT PRODUCT FORMATION CROSS SECTIONS AND TOTAL DIRECT REACTION CROSS SECTIONS

The cross section for the formation of light products in direct reactions are shown in table 1. For each ⁴⁰Ar energy two values of cross section are given: σ_{meas} and σ . The values of σ_{meas} have been obtained by integrating the differential cross sections $d\sigma(\vartheta)/d\Omega$ over the angle ϑ in the angular region where the measurements have been made. These angular intervals were determined using the solid curves drawn through the experimental points as shown in fig. 7. The values of σ were obtained by integrating the total angular distributions extrapolated to the region of small angles. The total cross sections of direct reactions for a ²³²Th+⁴⁰Ar were calculated from our data using the cross section for the formation of separate elements. These values should be considered as the lower limits of the total cross sections of direct processes. The character of the dependence of light product formation cross section on Z allows one to assume that the contribution to the total cross section from elements with charges Z larger and smaller than those shown in table 1 will be small. The total cross section is somewhat underestimated due mainly to the cutting-off of the Ar energy spectra.

4. Discussion of experimental results

4.1. ENERGY SPECTRA

In order to clarify the specific features of direct reactions with Ar ions it is useful to compare the kinetic energies of final products with the output Coulomb barrier height. Fig. 8 shows $(E_{c.m.}^{in} + Q_m - B_{c.m.}^{out})$ versus the angle of emission of the light products (c.m.s.). Here $E_{c.m.}^{in}$ is the initial kinetic energy, $B_{c.m.}^{out}$ is the output Coulomb barrier height (calculated for spherical nuclei having $r_0 = 1.46$ fm), Q_m is the Q-value of the reaction for the maximum product yield. The values $B_{c.m.}^{out}$ and Q_m were



Fig. 8. The difference of kinetic energies in the maximum energy distributions and Coulomb barrier heights of the final products with respect to the light product emission angle. ⁴⁰Ar ion energies in the middle plane of the target $E_0 = 288$ MeV and $E_0 = 379$ MeV.

calculated under the assumption of the two-body reaction and for the above isotopes. Note that these values are weakly dependent on the variations of the mass number A.

For few-nucleon transfers the curves $(E_{c.m.}^{in} + Q_m - B_{c.m.}^{out})$ have their maxima at angles somewhat smaller than the Rutherford scattering angle of ⁴⁰Ar ($\vartheta_{Ruth}^{e.m.} = 63^{\circ}$ for $E_0 = 288$ MeV and $\vartheta_{Ruth}^{e.m.} = 41^{\circ}$ for $E_0 = 379$ MeV). At these angles the kinetic energy of final products is close to $E_{c.m.}^{in}$ and the process determining the high-energy maximum in the energy spectrum is of quasi-elastic character. With decreasing angle, a sharp decrease in the maximum of the final kinetic energy spectrum takes place. It is reduced down to the value close to the height of the output Coulomb barrier, and is further weakly decreased with decreasing 9.

The low-energy maxima in the final kinetic energy spectra of products turn out to be lower than the output Coulomb barrier height and is smoothly decreased with increasing angle 9.

Kammuri¹⁸) when calculating the classical deflection of a heavy ion in the Coulomb and nuclear potentials has obtained a possible ion trajectory in traversing 0° with respect to the direction of the incident ion. In our case this suggests that the lowenergy maxima in the spectra of ions emitted at angles of 10° to 60° are indeed those emitted at angles of -10° to -60° . The points thus plotted are indicated by dashed lines in fig. 8 which shows that it is possible to connect both the low- and high-energy maxima in the energy spectra in one system.

For the multi-nucleon transfer reactions the kinetic energy of final products is somewhat smaller than the output Coulomb barrier height for spherical nuclei and varies smoothly with the angle ϑ . Note the similarity of the positions and the shapes of curves shown in fig. 8 for the utmost multi-nucleon transfers (O, F, Ne) and for the low-energy maxima of Cl, K, Ca. This fact may be considered as an indication that in both cases we may have the product trajectories traversing through 0° .

It is interesting that in some cases the energy spectrum maximum of a light product is below the output Coulomb barrier height (see figs. 6 and 8). Earlier the production of transfer reaction products at energies below the Coulomb barrier height has been observed on light and medium nuclei ^{19,20}). One can indicate some possible reasons for the formation of light products with such low energies: (i) the detected product is produced as a result of the decay of a deformed system with relatively smaller values of $B_{c.m.}^{out}$; (ii) the system decays into more than two fragments, of comparable charge and mass; (iii) the emission, by the excited light product, of α -particles and nucleons in the direction of motion. However, it is improbable that such a process as the decay of the system into more than two massive fragments determines the position of maxima in energy spectra. This is confirmed by the observation of the maxima of the energy spectra being near the Coulomb barrier for two particles in the outgoing channel and furthermore the noticeable limitation of energy spectra at relatively high energies.

There are no special arguments against the isotropic emission of nucleons or, which is more probable, of α -particles by a light product (after the decay of the system). With the isotropic decay of the light product the maximum in the energy spectrum will be determined by α -emission in the direction perpendicular to that of product motion. In this case the detection angle for a new product is changed slightly due to its great velocity. (The product energy in system decaying from the Coulomb barrier height is 4–5 MeV/nucleon.) At the same time, considering that the velocity of the light product after its decay is on an average retained, the variation of the kinetic energy of a new product with respect to the new Coulomb barrier is small. Indeed the points for O, F, Ne in fig. 8 are practically at the same curve. Thus, one may ²³²Th(⁴⁰Ar, X)

consider that at the moment of decay of the system, the nuclei are somewhat deformed and the deformation increases with the increase in the angle of the trajectory away from 0° .

It is interesting to note that the product yield in such processes is changed slightly over a wide range of atomic number $8 \le Z \le 20$ (see product yields in figs. 3-5 for the detection angle of 45° where such processes are separated sufficiently well).



Fig. 9. Partial angular distributions of light products with the fixed value of $Q = Q^*$. ⁴⁰Ar ion energy $E_0 = 288$ MeV. The curves for K with $Q^* = 0$ MeV and $Q^* = -20$ MeV have been calculated according to Strutinsky ²⁴). Calculation parameters are presented in table 2. The remaining curves have been drawn through the experimental points. All the points and curves for the given value of Q^* have been through the experimental points.

 Q^* have been multiplied by the coefficients given in the left-hand side of the figure.

4.2. ANGULAR DISTRIBUTIONS

In previous investigations ^{21, 22}) it has been shown that the shape of angular distributions of light products in transfer reactions can significantly change for the different parts of the energy spectrum. In this case the energy spectrum was separated into two parts and for each of them its own angular distribution was determined. Taking into account the large width of energy spectra it is reasonable in our case to

consider the angular distribution of reaction products at various excitation energies of the interacting system of two nuclei. Figs. 9 and 10 show the differential cross sections for the formation of reaction products as $d^2\sigma/dQd\Omega$ with the different Q-values, Q^* . The calculation was made under the assumption of the two particles in the outgoing channel. As is seen from these figures, in few-nucleon transfer reactions at small excitation energies of the final nuclei E^* ($E^* = Q_{gg} - Q^*$) the angular distribution





Fig. 10. Partial angular distributions of light products with the fixed value of $Q = Q^*$, ⁴⁰Ar ion energy $E_0 = 379$ MeV. Curves for K with $Q^* = 0$, $Q^* = -20$ MeV and $Q^* = -200$ MeV have been calculated according to Strutinsky 24). Calculation parameters are presented in table 2. The remaining curves are drawn through experimental points. All the points and curves for the given value

of Q^* have been multiplied by the coefficients shown in the bottom left-side of the figure.

has a pronounced maximum. With increasing excitation energy, the maximum in the angular distribution is shifted towards smaller angles, its width is increased, and a rise in the cross section at small angles appears. At higher excitation energies the yield of products increases monotonically towards 0°. The radius parameter r_0 calculated (for the closest approach of two nuclei) from the maximum positions in the angular distributions Cl and K with the minimum values of $|Q^*|$ is between (1.55–1.62)×10⁻¹³ cm.

Some investigations $^{18, 23-27}$ have been devoted to the theoretical analysis of the angular distributions of direct reaction products. Strutinsky's model 24 is most convenient for comparison with the experimental data. We have calculated the product angular distributions under the assumption of the "tangent" nuclear collisions and the quasi-elasticity of reactions. The function η_l determining reaction intensity in the channel with orbital momentum l has a maximum with some $l \approx l_0$ corresponding to the "tangent" trajectory. Taking

$$\eta_{l} = \eta_{l0} q(l - l_{0}) = \eta_{l0} q(x), \tag{1}$$

two expressions for the function q(x) are considered:

$$q(x) = \exp(-\alpha^2 x^2)$$
 (G), (2)

$$q(x) = \begin{cases} \exp(-\alpha x) & l > l_0 \\ 0 & l < l_0 \end{cases}$$
(E). (3)

The quantity $\alpha = 1/\Delta l$ determines the wave packet width.

It has been shown that the shape of the angular distribution for light reaction products depends upon the number of partial waves giving a contribution to the reaction and the classical deflection angle ϑ_{\max} . The maximum at the angle ϑ_{\max} appears only with the condition $\vartheta_{\max} \Delta l > 2$. Such an approach allows the extraction of the values Δl and ϑ_{\max} from the experimentally determined angular distributions.

TABLE 2The analysis of the angular distribution of the light product in the ²³²Th(⁴⁰Ar, K) reaction by
using the Strutinsky model ²⁴)

Energy ⁴⁰ Ar E ₀ (MeV)	Q* MeV	Calculation variant	ϑ _{max} (degrees)	<i>∆l</i> number of partial waves
288	0	G	57	12
	—20	G	56	9.2
379	0	G	36.5	25.1
	20	G	35.5	19.4
	200	E	5	6.3

Table 2 presents the results of calculations of wave packet widths for partial angular distributions of K with $Q^* = 0$ MeV and $Q^* = -20$ MeV. (Here the initial conditions of the model hold.) The calculated curves (variant G) are shown in figs. 9 and 10. The theoretical curves were normalized to the experimental cross sections at the angles ϑ_{max} . The agreement of the theoretical curves with the experimental points is quite satisfactory. In this new investigation Strutinsky has generalized his approach for inelastic direct processes 27). In this variant of the model the angular distribution of reaction products is determined by two factors: by quantum-mechanical dispersion due to the wave nature of the particle and by the dynamic angular dispersion due to nuclear interaction. The mathematical expressions for the angular distributions obtained here are equivalent to the expressions of ref. 24). But since the dynamic dispersion of a packet was not taken into account in the previous paper the parameter ΔI obtained from the analysis of the experimental angular distribution can be smaller than the real width of the wave packet.

However, it is impossible to distinguish the quantum-mechanical and dynamic dispersions within the framework of the proposed model ²⁷). We have made an attempt to compare the partial angular distributions shown in figs. 9 and 10 to those calculated using the formulas of the previous work ²⁴). It was found that for the largest $|Q^*|$ values with $E_0 = 379$ MeV the calculated angular distribution is consistent with the experimental, if the (E) variant is used. Fig. 10 shows the corresponding calculated curve for K with $Q^* = -200$ MeV. The calculated parameters are listed in table 2.

An attempt to fit the theoretical and experimental points for the intermediate values of Q^* using only the Gaussian or the exponent dependence of q(x) failed. Thus, for comparatively small $|Q^*|$ the character of the decrease in the angular distribution at the angles of $\vartheta > \vartheta_{max}$ requires definitely the application of (G), but then the observed increase of cross section at smaller angles cannot be explained on this basis. Perhaps it may be possible to explain the observed angular distributions by introducing a varying asymmetry in q(x), from (G) to (E), with increasing excitation energy.

4.3, DIRECT REACTION TOTAL CROSS SECTION AND THE CRITICAL ANGULAR MOMENTUM

The comparison of our data for the total cross section of direct reactions σ (see table 1) with the cross section for incomplete fusion σ_{ICF} obtained by the fission-fragment angular correlation experiments ²⁸) shows a noticeable divergence. The value σ in our experiments turned out to be considerably greater. It is possible that this divergence is due to the difficulties of separation, by the fission-fragment angular correlation method of the complete fusion processes and deep inelastic direct processes with large momentum transfers to the target nucleus.

The data obtained allow a determination of the value of the critical angular momentum $J_{\rm cr}$ for the ²³²Th+⁴⁰Ar system. Natowitz ²⁹) has remarked that the value $J_{\rm cr} = 170\hbar$ obtained from the data of Sikkeland ²⁸) may have been overestimated. In order to evaluate $J_{\rm cr}$ we have used the same formulas as in ref. ²⁹):

$$J_{\rm cr} = \left(\frac{\sigma_{\rm R} - \sigma_{\rm 1CF}}{\sigma_{\rm R}}\right)^{\frac{1}{2}} J_{\rm max}, \qquad (4)$$

 232 Th(40 Ar, X) 107

$$\sigma_{\rm R} = \pi R^2 \left(1 - \frac{B}{E_{\rm c.m.}} \right) \,, \tag{5}$$

$$J_{\rm max}^{2} = \frac{2\mu(E_{\rm c.m.} - B)R^{2}}{\hbar^{2}}.$$
 (6)

Here $\sigma_{\mathbf{R}}$ is the total reaction cross section, $\sigma_{\rm ICF}$ is the cross section for incomplete fusion, *B* is the input Coulomb barrier height, μ is the reduced mass, *R* is the sum of the target and projectile radii ($r_0 = 1.46$ fm), $J_{\rm max}$ is the maximum orbital angular momentum under the assumption of spherical nuclei with sharp boundaries. The results of calculation of $J_{\rm er}$ are presented in table 3. The values $J_{\rm cr}$ should be considered as the upper limit for the critical angular momentum, since the limiting value for $\sigma_{\rm ICF}$ was used in the calculations.

TABLE 3 Calculation of the critical angular momentum value of the 232 Th $+^{40}$ Ar system according to Natowitz 29)

Energy 40 Ar E ₀ (MeV)	σ _R (b)	σ _{ICF} (b)	J_{\max} (units of \hbar)	J_{cr} (units of \hbar) ^a)
288	1.9	1.1	157	102
379	2.9	2.4	222	94

^a) J_{cr} = the upper limit for the critical angular momentum value.

It should be noted that our data for J_{cr} values fit rather well the empirical systematics for critical angular momenta proposed by Natowitz²⁹).

5. Conclusion

The results obtained in this experiment demonstrate the possibility of the "element" approach in studying nuclear reactions with very heavy ions.

Transfer reactions with ⁴⁰Ar have been shown to occur both in the form of quasielastic and deep inelastic processes. The former are characteristic of few-nucleon transfer reactions. In this case the reaction product retains a major part of the incident ion velocity, whereas the product angular distribution has a pronounced peak at angles close to the Rutherford scattering angle for the incident ion. Deep inelastic processes make a noticeable contribution to few-nucleon transfer reactions and are dominant in multi-nucleon reactions. In such processes all the kinetic energy of nuclear collisions is spent in the rearrangement and excitation of nuclei. The angular distributions are described by a monotonic increase of the yield with decreasing detection angle.

Such processes are an intermediate stage to the production of the compound system. This may be suggested by the transition of the trajectory of light product throughout 0° and deformed configuration of the system at the moment of scission.

The transfer of a considerable number of nucleons from a heavy ion to the target

nucleus with a noticeable cross section indicates the possibility of using such processes for the synthesis of transuranic elements and, possibly, of superheavy nuclei in a new region of stability. It is worth noting that the large width of the energy spectra of light products increases the probability of processes resulting in the weak excitation of the final heavy products.

The authors are deeply indebted to Academician G. N. Flerov for his interest to this investigation and valuable discussions and to B. A. Zager and the cyclotron operation staff for their cooperation.

References

- 1) Yu. Ts. Oganessjan, Yu. E. Penionzhkevich and O. A. Shamsutdinov, Yad. Fiz. 14 (1971) 54
- 2) A. G. Demin, V. Kusch, M. B. Miller, A. S. Pasyuk, A. A. Pleve and Yu. P. Tretiakov, Proc. of the Int. Conf. on heavy-ion physics, Dubna, 11–17 Febr. 1971, JINR, D7–5769 (Dubna, 1971) p. 169
- 3) R. Bimbot, C. Deprun, H. Gauvin, Y. L. Beyec, M. Lefort, J. Peter and B. Tamain, Nature 234 (1972) 215
- 4) G. N. Flerov, Yu. Ts. Oganessjan, JINR, P7-6523, Dubna, 1972
- 5) A. G. Artukh, V. V. Avdeichikov, G. F. Gridnev, V. L. Mikheev, V. V. Volkov and J. Wilczynski, Nucl. Phys. A176 (1971) 284
- 6) H. Kumpf and E. D. Donets, JETP (Sov. Phys.) 17 (1963) 539
- W. Grochulski, T. Kwiecinska, Lian Go-chan, E. Lozynski, J. Maly, L. K. Tarasov and V. V. Volkov, Proc. Third Conf. on reactions between complex nuclei, ed. A. Chiorso, R. M. Diamond, H. E. Conzett (University of California Press, Berkeley, 1963) p. 120
- 8) E. Lozynski and I. I. Chuburkova, JINR, P7-3630, Dubna, 1967
- 9) A. G. Artukh, V. V. Avdeichikov, J. Ero, G. F. Gridnev, V. L. Mikheev and V. V. Volkov, Nucl. Instr. 83 (1970) 72
- 10) M. W. Sachs, C. Chasman and D. A. Bromley, Nucl. Instr. 41 (1966) 213
- 11) G. W. Butler, A. M. Poskanzer and D. A. Landis, Nucl. Instr. 89 (1970) 189
- 12) A. G. Artukh, V. V. Volkov, L. Pomorski and J. Tys, JINR, P7-5494, Dubna, 1970
- 13) L. C. Northcliffe and R. F. Schilling, Nucl. Data Tables A7 (1970) 233
- 14) A. G. Artukh, J. Wilczynski, V. V. Volkov, G. F. Gridnev and V. L. Mikheev, JINR, P7-6815, Dubna, 1973
- S. A. Karamjan, Yu. Ts. Oganessjan, Yu. E. Penionzhkevich and B. J. Pustilnik, JINR, P7-5884, Dubna, 1971
- 16) K. Alder, A. Bohr, T. Huns, B. Mottelson and A. Winter, Rev. Mod. Phys. 28 (1956) 432
- 17) E. Lozynski, JINR, P-1486, Dubna, 1963
- 18) T. Kammuri, Progr. Theor. Phys. 28 (1962) 934
- 19) J. Wilczynski, V. V. Volkov and P. Decowski, Yad. Fiz. 5 (1967) 942
- 20) G. F. Gridnev, V. V. Volkov and J. Wilczynski, Nucl. Phys. A142 (1970) 385
- 21) A. G. Artukh, G. F. Gridnev, V. L. Mikheev, J. Wilczynski, V. V. Volkov, Proc. of the Int. Conf. on nuclear reactions induced by heavy ions, ed. R. Bock and W. R. Hering (North-Holland, Amsterdam, 1970) p. 140
- 22) J. Galin, D. Guerreay, M. Lefort, J. Peter, X. Tarrago and R. Basile, Nucl. Phys. A159 (1970) 461
- 23) B. N. Kalinkin, J. Grabowski, Proc. Third Conf. on reactions between complex nuclei, Asilomar, 1963 (Univ. Calif. Press, 1963) p. 129
- 24) V. M. Strutinsky, ZhETF (USSR) 46 (1964) 2078
- 25) W. E. Fraha and R. H. Venter, Nucl. Phys. 59 (1964) 651
- 26) R. da Silveira, Communications European Conf. on nucl. physics, Aix-en-Provence (France), 1972 (Compt. Rend., 1972) p. 57
- 27) V. M. Strutinsky, Phys. Lett. 44B (1973) 245
- 28) T. Sikkeland, UCRL-16580 (1965)
- 29) J. B. Natowitz, Phys. Rev. C1 (1970) 623

108